









•In most studies of treatment conducted over the last 40 years, the average treated person is better off than 80% of the untreated sample.

- •The outcome of behavioral health services equals and, in most cases, exceeds medical treatments.
- •On average, mental health professionals achieve outcomes on par with success rates obtained in randomized clinical trials (with and without co-morbidity).

Duncan, B., Miller, S., Wampold, B., & Hubble, M. (eds.) (2009). The Heart and Soul of Change: Delivering What Works. Washington, D.C.: APA Press.

Minami, T., Wampold, B., Serlin, R., Hamilton, E., Brown, G., Kircher, J. (2008). Benchmarking for psychotherapy efficacy. *Journal of Consulting and Clinical Psychology*, 75 232-243.



The Evidence:

Three "Stubborn" Facts

- Drop out rates average 47%;
- •Mental health professionals frequently fail to identify failing cases;
- •1 out of 10 consumers accounts for 60-70% of expenditures.

Aubrey, R., Self, R., & Halstead, J. (2003). Early non attendance as a predictor of continued non-attendance and subsequent attribtion from psychological help. Clinical Psychology, 32, 6-10.

Chasson, G. (2005). Attrition in child treatment. Psychotherapy Bulletin, 40(1), 4-7.

Harmon, S.J., Lambert, M.J., Smart, D.M., Hawkins, E., Nielsen, S.L., Slade, K., Lutz, W., (2007) Enhancing outcome for potential treatment failures: Therapist-client feedback and clinical support tools. *Psychotherapy Research*, 17(4), 379-392


Lambert, M.J., Whipple, J., Hawkins, E., Vermeersch, D., Nielsen, S., & Smart, D. (2004). Is it time for clinicians routinely to track client outcome? A meta-analysis. *Clinical Psychology*, *10*, 288-301.






















The Evidence

•FIT is being used with broad and diverse group of adults, youth, and children in agencies and treatment settings around the world including:

- Inpatient
- Outpatient
- •Residential
- •Prison-based (mandated care)
- •Case management





Bohanske, B. & Franczak, M. (2009). Transforming public behavioral health care: A case example of consumer directed services, recovery, and the common factors. In B. Duncan, S. Miller, B. Wampold, & M. Hubble. (Eds.) (2009). *The Heart and Soul of Change* (2nd Ed.). Washington, D.C.: APA Press.

What Works in Therapy

Consumers:	Clinicians:	Payers:
Individualized care	Professional autonomy	Accountability
Needs met in the most effective and efficient manner possible (value-based purchasing)	Ability to tailor treatment to the individual client(s) and local norms	Efficient use of resources
Ability to make an informed choice regarding treatment providers	Elimination of invasive authorization and oversight procedures	Better relationships with providers and decreased management costs
A continuum of possibilities for meeting care needs	Paperwork and standards that facilitate rather than impede clinical work	Documented return on investment

FIT Fits

•In the Task Force's recent report (APA, 2006), the following definition for EBPP was set forth: "Evidence-based practice in psychology (EBPP) is the integration of the best available research with clinical expertise in the context of patient characteristics, culture, and preferences" (p. 273; emphasis included in the original text). Regarding the phrase "clinical expertise" in this definition, the Task Force expounded the following (APA, 2006; p. 276-277).

•Clinical expertise also entails the monitoring of patient progress (and of changes in the patient's circumstances—e.g.,job loss, major illness) that may suggest the need to adjust the treatment (Lambert, Bergin, & Garfield,2004a). If progress is not proceeding adequately, the psychologist alters or addresses problematic aspects of the treatment (e.g., problems in the therapeutic relationship or in the implementation of the goals of the treatment) as appropriate.

*COTTIMULA (


Presidential task force on evidence-based practice. (2006). Evidence-based practice in psychology. *American Psychologist*, 61(4), 271-285.








Three Steps for becoming FIT:


- Create a "Culture of feedback";
- 2. Integrate alliance and outcome feedback into clinical care;
- 3. Learn to "fail successfully."











Step One:


Creating a "Culture of Feedback"


Outcome Rating Scale (ORS)


Name Age (Yrs):
Session # Date:


Looking back over the last week, including today, help us understand how you have been feeling by rating how well you have been doing in the following areas of your life, where marks to the left represent low levels and marks to the right indicate high levels.


- •When scheduling a first appointment, provide a rationale for seeking client feedback regarding outcome:
 - Work a little differently;
 - •If we are going to be helpful should see signs sooner rather than later;
 - •If our work helps, can continue as long as you like;
 - •If our work is not helpful, we'll seek consultation (session 3 or 4), and consider a referral (within no later than 8 to 10 visits).


























Step Two:

Integrating Feedback into Care

•Computer-generated "trajectories of change":


- •Uses a normative database and linear regression to plot clientspecific trajectories;
- •Depicts the amount of change in scores needed to be attributable to treatment.



Step Two:

Integrating Feedback into Care

"Therapists typically are not cognizant of the trajectory of change of patients seen by therapists in general... That is to say, they have no way of comparing their treatment outcomes with those obtained by other therapists."





Wampold, B., & Brown, J. (2006). Estimating variability in outcomes attributable to therapists: A naturalistic study of outcomes in managed care. *Journal of Consulting and Clinical Psychology*, 73 (5), 914-923.



